

WHITE FLINT PHARMACY
11125 Rockville Pike , Rockville , MD , 20852
Phone#: 3018813828 Fax#: 3017709629

Drug NDC: 00525-6748-01
Drug Name: PALGIC

What is this medicine?

CARBINOXAMINE (kar bi NOX a meen) is an antihistamine. It is used to treat the symptoms of allergies and hives.

This medicine may be used for other purposes; ask your health care provider or pharmacist if you have questions.

How should I use this medicine?

Take this medicine by mouth with a glass of water. Follow the directions on the prescription label. Take your medicine at regular intervals. Do not take it more often than directed. Talk to your pediatrician regarding the use of this medicine in children. Special care may be needed.

Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once. NOTE: This medicine is only for you. Do not share this medicine with others.

What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions: asthma, glaucoma, heart disease, high blood pressure, pain or difficulty passing urine, stomach problems, taking MAOIs like Carbox, Eldepryl, Marplan, Nardil, and Parnate, an unusual or allergic reaction to carbinoxamine, other medicines, foods, dyes, or preservatives like sulfites, pregnant or trying to get pregnant, breast-feeding

What if I miss a dose?

If you miss a dose, take it as soon as you can. If it is almost time for your next dose, take only that dose. Do not take double or extra doses.

Where should I keep my medicine?

Keep out of the reach of children. Store at room temperature between 15 and 30 degrees C (59 and 86 degrees F). Keep medicine bottle tightly closed. Protect from light. Throw away any unused medicine after the expiration date.

NOTE: This sheet is a summary. It may not cover all possible information. If you have questions about this medicine, talk to your doctor, pharmacist, or health care provider.

What should I watch for while using this medicine?

Tell your doctor or healthcare professional if your symptoms do not start to get better or if they get worse. You may get drowsy or dizzy. Do not drive, use machinery, or do anything that needs mental alertness until you know how this medicine affects you. Do not stand or sit up quickly, especially if you are an older patient. This reduces the risk of dizzy or fainting spells. Alcohol may interfere with the effect of this medicine. Avoid alcoholic drinks. Your mouth may get dry. Chewing sugarless gum or sucking hard candy, and drinking plenty of water may help. Contact your doctor if the problem does not go away or is severe. This medicine may cause dry eyes and blurred vision. If you wear contact lenses you may feel some discomfort. Lubricating drops may help. See your eye doctor if the problem does not go away or is severe.

What side effects may I notice from receiving this medicine?

Side effects that you should report to your doctor or health care professional as soon as possible: agitation, nervousness, excitability, not able to sleep, blurred vision, fainting spells, fast, irregular heartbeat, fever, difficulty passing urine, seizures, slurred speech, confusion, tremor. Side effects that usually do not require medical attention (report to your doctor or health care professional if they continue or are bothersome): constipation or diarrhea, headache, loss of appetite, nausea, vomiting, stomach upset

This list may not describe all possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.