

**WHITE FLINT PHARMACY**  
**11125 Rockville Pike , Rockville , MD , 20852**  
**Phone#: 3018813828 Fax#: 3017709629**

**Drug NDC: 43386-0500-11**  
**Drug Name: FAMOTIDINE**

**What is this medicine?**

FAMOTIDINE (fa MOE ti deen) is a type of antihistamine that blocks the release of stomach acid. It is used to treat stomach or intestinal ulcers. It can also relieve heartburn from acid reflux.

**This medicine may be used for other purposes; ask your health care provider or pharmacist if you have questions.**

**How should I use this medicine?**

Take this medicine by mouth. Follow the directions on the prescription label. Shake the bottle for 5 to 10 seconds. Use a specially marked spoon or container to measure your medicine. Ask your pharmacist if you do not have one. Household spoons are not accurate. If you only take this medicine once a day, take it at bedtime. Take your doses at regular intervals. Do not take your medicine more often than directed. Talk to your pediatrician regarding the use of this medicine in children. Special care may be needed.

**Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once. NOTE: This medicine is only for you. Do not share this medicine with others.**

**What should I tell my health care provider before I take this medicine?**

They need to know if you have any of these conditions: kidney or liver disease an unusual or allergic reaction to famotidine, other medicines, foods, dyes, or preservatives pregnant or trying to get pregnant breast-feeding

**What if I miss a dose?**

If you miss a dose, take it as soon as you can. If it is almost time for your next dose, take only that dose. Do not take double or extra doses.

**Where should I keep my medicine?**

Keep out of the reach of children. Store at room temperature between 15 and 30 degrees C (59 and 86 degrees F). Do not freeze. Throw away any unused suspension after thirty days.

**NOTE: This sheet is a summary. It may not cover all possible information. If you have questions about this medicine, talk to your doctor, pharmacist, or health care provider.**

**What should I watch for while using this medicine?**

Tell your doctor or health care professional if your condition does not start to get better or if it gets worse. Finish the full course of medicine prescribed, even if you feel better. Do not take with aspirin, ibuprofen or other anti-inflammatory medicines. These can make your condition worse. Do not smoke cigarettes or drink alcohol. These cause irritation in your stomach and can increase the time it will take for ulcers to heal. If you get black, tarry stools or vomit up what looks like coffee grounds, call your doctor or health care professional at once. You may have a bleeding ulcer.

**What side effects may I notice from receiving this medicine?**

Side effects that you should report to your doctor or health care professional as soon as possible: agitation, nervousness, confusion, hallucinations, skin rash, itching. Side effects that usually do not require medical attention (report to your doctor or health care professional if they continue or are bothersome): constipation, diarrhea, dizziness, headache

**This list may not describe all possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.**