WHITE FLINT PHARMACY

11125 Rockville Pike , Rockville , MD , 20852

Drug NDC: 55390-0337-01
Drug Name: DACTINOMYCIN

What is this medicine?

DACTINOMYCIN (dak ti noe MYE sin) is a chemotherapy drug. It is used to treat many kinds of cancer like Wilms' tumor, some sarcomas, and placental and testicular cancers. It is also used to treat other solid tumors.

This medicine may be used for other purposes; ask your health care provider or pharmacist if you have questions.

How should I use this medicine?

This drug is given as an infusion into a vein. It is administered in a hospital or clinic by a specially trained health care professional.\n\nTalk to your pediatrician regarding the use of this medicine in children. While this drug may be prescribed for children as young as 6 months of age for selected conditions, precautions do apply.

Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once.\nNOTE: This medicine is only for you. Do not share this medicine with others.

What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions:\n-infection (especially a virus infection such as chickenpox, cold sores, or herpes)\n-liver disease\n-low blood counts like low platelets, red blood cells, white blood cells\n-recent radiation therapy\n-an unusual or allergic reaction to dactinomycin, other chemotherapy agents, other medicines, foods, dyes, or preservatives\n-pregnant or trying to get pregnant\n-breast-feeding

What if I miss a dose?

It is important not to miss your dose. Call your doctor or health care professional if you are unable to keep an appointment.

Where should I keep my medicine?

This drug is given in a hospital or clinic and will not be stored at home.

NOTE: This sheet is a summary. It may not cover all possible information. If you have questions about this medicine, talk to your doctor, pharmacist, or health care provider.

What should I watch for while using this medicine?

Your condition will be monitored carefully while you are receiving this medicine. You will need important blood work done while you are taking this medicine. In InThis drug may make you feel generally unwell. This is not uncommon, as chemotherapy can affect healthy cells as well as cancer cells. Report any side effects. Continue your course of treatment even though you feel ill unless your doctor tells you to stop. In InCall your doctor or health care professional for advice if you get a fever, chills or sore throat, or other symptoms of a cold or flu. Do not treat yourself. This drug decreases your body's ability to fight infections. Try to avoid being around people who are sick. In InThis medicine may increase your risk to bruise or bleed. Call your doctor or health care professional if you notice any unusual bleeding. In InTalk to your doctor about your risk of cancer. You may be more at risk for certain types of cancers if you take this medicine. In InDo not become pregnant while taking this medicine. Women should inform their doctor if they wish to become pregnant or think they might be prognant. There is a potential for serious side effects to an unborn child. Talk to your health care professional or pharmacist for more information. Do not breast-feed an infant while taking this medicine.

What side effects may I notice from receiving this medicine?

Side effects that you should report to your doctor or health care professional as soon as possible:\n-allergic reactions like skin rash, itching or hives, swelling of the face, lips, or tongue\n-low blood counts - this medicine may decrease the number of white blood cells, red blood cells and platelets. You may be at increased risk for infections and bleeding.\n-signs of infection - fever or chills, cough, sore throat, pain or difficulty passing urine\n-signs of decreased platelets or bleeding - bruising, pinpoint red spots on the skin, black, tarry stools, blood in the urine\n-signs of decreased red blood cells - unusually weak or tired, fainting spells, lightheadedness\n-breathing problems\n-changes in vision\n-chest pain\n-mount sores\n-pain, swelling, redness at site where injected\n-seizures\n-stomach pain, ulcers\n-swelling of the ankles, feet, hands, or stomach\n-trouble passing urine or change in the amount of urine\n-vomiting\n-y-yellowing of the eyes or skin\n\n\Side effects that usually do not require medical attention (report to your doctor or health care professional if they continue or are bothersome)\n-acne-like skin rash\n-darker skin color\n-diarrhea\n-hair loss\n-loss of appetite\n-nausea

This list may not describe all possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.