

WHITE FLINT PHARMACY
11125 Rockville Pike , Rockville , MD , 20852
Phone#: 3018813828 Fax#: 3017709629

Drug NDC: 64661-0032-30
Drug Name: J-TAN D PD

What is this medicine?

BROMPHENIRAMINE; PSEUDOEPHEDRINE (brome fen IR a meen; soo doe e FED rin) is a combination of an antihistamine and a decongestant. It is used to treat the symptoms of allergy and colds. This medicine will not treat an infection.

This medicine may be used for other purposes; ask your health care provider or pharmacist if you have questions.

How should I use this medicine?

Take this medicine by mouth. Follow the directions on the prescription label. Use a specially marked spoon or container to measure each dose. Ask your pharmacist if you do not have one. Household spoons are not accurate. You can take it with or without food. If it upsets your stomach, take it with food. Take your medicine at regular intervals. Do not take your medicine more often than directed.
Talk to your pediatrician regarding the use of this medicine in children. While this drug may be prescribed for selected conditions, precautions do apply.
Patients over 65 years old may have a stronger reaction and need a smaller dose.

Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once.
NOTE: This medicine is only for you. Do not share this medicine with others.

What should I tell my health care provider before I take this medicine?

They need to know if you have any of the following conditions:
asthma
diabetes
difficulty passing urine
glaucoma
heart disease
high blood pressure
taken an MAOI like Carbox, Eldepryl, Marplan, Nardil, or Parnate in last 14 days
thyroid disease
ulcer or other stomach problem
an unusual or allergic reaction to brompheniramine, pseudoephedrine, other medicines, foods, dyes, or preservatives
pregnant or trying to get pregnant
breast-feeding

What if I miss a dose?

If you miss a dose, take it as soon as you can. If it is almost time for your next dose, take only that dose. Do not take double or extra doses.

Where should I keep my medicine?

Keep out of the reach of children.
Store at room temperature between 20 and 25 degrees C (68 and 77 degrees F). Protect from light. Throw away any unused medicine after the expiration date.

NOTE: This sheet is a summary. It may not cover all possible information. If you have questions about this medicine, talk to your doctor, pharmacist, or health care provider.

What should I watch for while using this medicine?

Tell your doctor or healthcare professional if your symptoms do not start to get better or if they get worse. If you have trouble falling asleep at night, take the last dose of the day at least a few hours before bedtime.
You may get drowsy or dizzy. Do not drive, use machinery, or do anything that needs mental alertness until you know how this medicine affects you. To reduce the risk of dizzy or fainting spells, do not stand or sit up quickly, especially if you are an older patient. Alcohol may increase dizziness and drowsiness. Avoid alcoholic drinks.
Your mouth may get dry. Chewing sugarless gum or sucking hard candy, and drinking plenty of water may help. Contact your doctor if the problem does not go away or is severe.

What side effects may I notice from receiving this medicine?

Side effects that you should report to your doctor or health care professional as soon as possible:
allergic reactions like skin rash, itching or hives, swelling of the face, lips, or tongue
breathing problems
confusion
fast or irregular heartbeat
feeling faint or lightheaded, falls
hallucinations
high blood pressure
seizure
trouble passing urine or change in the amount of urine
vomiting
Side effects that usually do not require medical attention (report to your doctor or health care professional if they continue or are bothersome):
anxiety, excitement, irritability, nervousness
diarrhea
headache
loss of appetite
stomach upset, nausea

This list may not describe all possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.